

Asymptotic Padé Approximants and the SQCD β -function

I. Jack, D.R.T. Jones

Dept of Mathematical Sciences, University of Liverpool, Liverpool L69 3BX, UK

M.A. Samuel

Department of Physics, Oklahoma State University, Stillwater, Oklahoma 74078, USA

We present a prediction for the four loop β -function for SQCD based on the method of Asymptotic Padé Approximants.

June 1997

Recently a four-loop calculation of the gauge β -function (β_g) in an $N = 1$ supersymmetric gauge theory was presented[1]. The computation was performed within the usual supersymmetric dimensional regularisation scheme (DRED), which is standard for perturbative calculations in supersymmetry and which consequently would be the most convenient for any future phenomenological applications. The result was derived somewhat indirectly, by starting from a partial calculation for the abelian case. The coupling constant redefinition relating the abelian result to the corresponding exact NSVZ result [2] could then be constructed, and this redefinition was then extended to the non-abelian case by imposing the vanishing of the β -function beyond one loop in the case of $\mathcal{N} = 2$ supersymmetry. However, one possible redefinition which was present in general, but vanished for the abelian case, could not be determined, and consequently the four-loop result β_3 contained one undetermined constant. Our purpose here is to predict the value of this constant by using improved Padé approximation techniques which have recently been developed and applied to the case of standard QCD[3]. In that case a prediction for β_3 was found to be in good agreement with a subsequent analytic calculation[4], especially when the contributions involving quartic Casimir group theory structures (which arise for the first time at four loops) are omitted in making the comparison. In fact these structures simply do not arise in the supersymmetric case which concerns us here. This observation, allied with the fact that we have more information about β_3 than was available to Ellis et al in the QCD case, makes us optimistic regarding the accuracy of our prediction.

The method is as follows: For a perturbative series $P(x) = \sum_{n=0}^{\infty} S_n x^n$, the Padé approximant $P_{[N/M]}(x)$ is a ratio of polynomials $A_N(x)$ and $B_M(x)$, of degree N and M respectively, chosen so that

$$P_{[N/M]} = \frac{A_N(x)}{B_M(x)} = P(x) + O(x^{N+M+1}). \quad (1)$$

It can be argued that in the case of QCD (and presumably also of supersymmetric QCD) the relative error

$$\delta_{N+M+1} \equiv \frac{S_{N+M+1}^{\text{Padé}} - S_{N+M+1}}{S_{N+M+1}} \quad (2)$$

has the asymptotic form

$$\delta_{N+M+1} \simeq -\frac{M!A^M}{L^M} \quad (3)$$

as $N \rightarrow \infty$, for fixed M , where A is a constant and $L = N + M + aM + b$. We will be concerned with $[0, 1]$ and $[1, 1]$ Padés, so $M = 1$; and we choose[3] a, b so that $a + b = 0$. The Asymptotic-Padé Approximant Prediction (APAP) is then given by:

$$S_{N+M+1}^{\text{APAP}} = \frac{S_{N+M+1}^{\text{Pade}}}{1 + \delta_{N+M+1}}. \quad (4)$$

If S_0, S_1 and S_2 are known, then the prediction for S_3 is obtained as follows. Firstly by matching to $P(x)$ up to linear terms as in Eq. (1), we obtain

$$P_{[0,1]} = \frac{S_0^2}{S_0 - S_1 x}, \quad (5)$$

giving as a prediction for the coefficient of the x^2 term $S_2^{\text{Pade}} = \frac{S_1^2}{S_0}$. Then from Eq. (2) we find $\delta_2 = \frac{S_1^2}{S_0 S_2} - 1$. From Eq. (3) we deduce $A = -\delta_2$, and thence $\delta_3 = -\frac{1}{2}A = \frac{1}{2}\delta_2$. The $[1, 1]$ Padé prediction for S_3 is easily derived to be $S_3^{\text{Pade}} = \frac{S_2^2}{S_1}$, and we then use Eq. (4) to give an improved estimate S_3^{APAP} incorporating δ_3 .

The gauge β -function is given by

$$\beta_g = g \sum_{n=0}^{\infty} \beta_n x^{n+1}, \quad (6)$$

where $x = \frac{g^2}{16\pi^2}$, and β_n corresponds to the $(n+1)$ -loop result in perturbation theory. The procedure given above yields a prediction for the four-loop contribution β_3 . The final feature of the method is to perform the above process for a range of values of N_f , the number of flavours, and then to match the results to a cubic polynomial in N_f . In the QCD case this enabled the authors of Ref. [3] to incorporate as an extra piece of information the known coefficient of N_f^3 , obtained by large- N_f calculations. In our present case we shall be able to exploit the fact that we know the form of β_3 as a function of N_f up to a single parameter. The results of Ref. [1] were presented for a general theory; for the present purposes, however, we specialise to supersymmetric QCD, obtaining [5]:

$$\beta_0 = N_f - 3N_c \quad (7a)$$

$$\beta_1 = \left[4N_c - \frac{2}{N_c} \right] N_f - 6N_c^2 \quad (7b)$$

$$\beta_2 = \left[\frac{3}{N_c} - 4N_c \right] N_f^2 + \left[21N_c^2 - \frac{2}{N_c^2} - 9 \right] N_f - 21N_c^3 \quad (7c)$$

$$\beta_3 = \mathcal{A} + \mathcal{B}N_f + \mathcal{C}N_f^2 + \mathcal{D}N_f^3 \quad (7d)$$

where N_c is the number of colours, and

$$\begin{aligned}
\mathcal{A} &= -(6 + 36\alpha)N_c^4 \\
\mathcal{B} &= 36(1 + \alpha)N_c^3 - (34 + 12\alpha)N_c - \frac{8}{N_c} - \frac{4}{N_c^3} \\
\mathcal{C} &= -\left(\frac{62}{3} + 2\kappa + 8\alpha\right)N_c^2 + \frac{100}{3} + 4\alpha + \frac{6\kappa - 20}{3N_c^2} \\
\mathcal{D} &= -\frac{2}{3N_c}.
\end{aligned} \tag{8}$$

Here $\kappa = 6\zeta(3)$ and α is the constant which we hope to determine¹.

Note that $\mathcal{D}$ does not involve α . It is the leading- N_f contribution at four loops and could, of course, have been obtained from the large- N_f results presented in Refs. [5], [7]. We may therefore proceed in much the same way as did the authors of Ref. [3]. We first compute S_3^{APAP} as described above, and then after determining $\mathcal{A}, \mathcal{B}, \mathcal{C}$ from the fit to Eq. (7d) we can obtain three distinct “predictions” for α , $\alpha(\mathcal{A})$, $\alpha(\mathcal{B})$ and $\alpha(\mathcal{C})$. Naturally a test of the approach is the extent to which these predictions agree and are insensitive to the input value of N_c . As in Ref. [3] we use input values $N_f = 0, 1, 2, 3, 4$, and used a value for A , $\langle A \rangle$, obtained by averaging $A(N_f)$ for the input values; the outcome is not very sensitive, in fact, to whether we use $\langle A \rangle$ or $A(N_f)$. The results of this procedure are shown in Table 1:

N_c	$\alpha(\mathcal{A})$	$\alpha(\mathcal{B})$	$\alpha(\mathcal{C})$	$\alpha(\mathcal{C}) (\kappa = 0)$
2	2.56	2.34	-0.31	1.62
3	2.43	2.42	0.50	2.38
4	2.46	2.47	0.63	2.48
5	2.45	2.46	0.74	2.57

Table 1: APAP predictions for the unknown parameter α .

The results for $\alpha(\mathcal{A}), \alpha(\mathcal{B})$ are remarkable for their consistency and stability. In the fourth column of the table we show the result for $\alpha(\mathcal{C})$ consequent on omitting from β_3

¹ It is also possible to apply the APAP method to the cases of QED or supersymmetric QED, for which the complete four-loop results can be extracted from Refs. [6] or [1] respectively. However, the form of the series in these cases is not conducive to the accuracy of the approximation and the four-loop predictions are somewhat less impressive. We hope to discuss these and other applications of the APAP method elsewhere.

the terms proportional to κ . It could be argued that this is natural since, given that $\zeta(3)$ occurs for the first time at four loops, these terms cannot be accurately produced by the Padé. (This is similar to the argument made with respect to the quartic Casimir terms in Ref. [3].) In any case, our results clearly suggest a value for α of around 2.45.

An alternative approach is as follows. The unknown parameter α in Eq. (8) is one of three parameters that were introduced in Ref. [1], each being the coefficient of a certain coupling constant redefinition. Two of these parameters were in fact determined in Ref. [1]; but we can test the reliability of the APAP method by reintroducing one of them and comparing the APAP prediction with a β_3 now dependent on two parameters. Thus we replace β_3 by $\tilde{\beta}_3$ where:

$$\beta_3 \rightarrow \tilde{\beta}_3 = \tilde{\mathcal{A}} + \tilde{\mathcal{B}}N_f + \tilde{\mathcal{C}}N_f^2 + \tilde{\mathcal{D}}N_f^3 \quad (9)$$

and

$$\begin{aligned} \tilde{\mathcal{A}} &= \mathcal{A}, \\ \tilde{\mathcal{B}} &= \mathcal{B} + \delta \left[24N_c^3 - 12 \left(N_c + \frac{1}{N_c} \right) \right] \\ \tilde{\mathcal{C}} &= \mathcal{C} + \delta \left[-20N_c^2 + 16 + \frac{4}{N_c^2} \right] \\ \tilde{\mathcal{D}} &= \mathcal{D} + \delta \left[4N_c - \frac{4}{N_c} \right]. \end{aligned} \quad (10)$$

In the notation of Ref. [1], $\alpha \equiv \alpha_1$ and $\delta \equiv -\frac{2}{3} - \alpha_2$. In Ref. [1] it was shown that $\alpha_2 = -\frac{2}{3}$, and so we hope to find that $\delta = 0$. For fixed α we determine three distinct “predictions” for δ by fitting S_3^{APAP} to a polynomial as before. As an example, for $N_c = 4$ we obtain:

$$\delta(\tilde{\mathcal{B}}) = 7.85\alpha - 19.24 \quad (11a)$$

$$\delta(\tilde{\mathcal{C}}) = 19.38\alpha - 48.16 \quad (11b)$$

$$\delta(\tilde{\mathcal{D}}) = 51.94\alpha - 126.87. \quad (11c)$$

If we omit κ terms then Eqs.(11a), (11c) are unaffected, but Eq.(11b) becomes

$$\delta(\tilde{\mathcal{C}}) = 19.38\alpha - 47.41.$$

When we plot $\delta(\tilde{\mathcal{B}})$, $\delta(\tilde{\mathcal{C}})$ and $\delta(\tilde{\mathcal{D}})$ against α , as in Fig. 1, the results are quite striking. We see that $\delta(\tilde{\mathcal{B}})$, $\delta(\tilde{\mathcal{C}})$ and $\delta(\tilde{\mathcal{D}})$ are zero for very similar values of α , around 2.44–2.49. So the predicted values for $\delta(\tilde{\mathcal{B}})$, $\delta(\tilde{\mathcal{C}})$ and $\delta(\tilde{\mathcal{D}})$ agree rather well with the correct value $\delta = 0$ if $2.44 \leq \alpha \leq 2.49$, and this may be interpreted as a prediction for α .




Fig. 1: Graph of $\delta(\tilde{\mathcal{B}})$ (solid line), $\delta(\tilde{\mathcal{C}})$ (dashed line) and $\delta(\tilde{\mathcal{D}})$ (dash-dotted line) against α for $N_c = 4$.




Fig. 2: Graph of $\delta(\tilde{\mathcal{B}})$, $\delta(\tilde{\mathcal{C}})$ and $\delta(\tilde{\mathcal{D}})$ against α for $N_c = 4$ (omitting κ terms).

The convergence is again even better if we omit the term in κ , as depicted in Fig. 2, where each δ crosses the axis at $\alpha = 2.44$. As N_c is increased, we find that the convergence of the δ s improves in the case where we retain κ ; in fact for large N_c all the δ s are zero for $\alpha = 2.43$ irrespective of whether or not we retain κ .

What is our final prediction for α ? The relatively simple relationship between the NSVZ β -function and the corresponding DRED one as explored in Ref. [1] suggests that α is a simple fraction; given our results here, then we would expect $\alpha = 12/5$ or perhaps $\alpha = 5/2$. ($\alpha = 17/7$ is even closer, but experience suggests that this number is unlikely to emerge from a perturbative calculation.) The result $\alpha = 2\zeta(3)$ is also possible; in which case the apparent better convergence produced by excising κ from the comparison would need to be dismissed as coincidental.

Acknowledgements

This work arose out of discussions at the “Beyond the Standard Model V” Conference at Balholm, Norway. We thank Per Osland and the other organisers of the meeting for providing a pleasant and stimulating atmosphere. MAS was supported by the US Department of Energy under Grant No. DE-FG05-84ER40215.

References

- [1] I. Jack, D.R.T. Jones and C.G. North, Nucl. Phys. **B486** (1997) 479
- [2] V. Novikov et al, Nucl. Phys. **B229** (1983) 381;
V. Novikov et al, Phys. Lett. **B166** (1986) 329;
M. Shifman and A. Vainshtein, Nucl. Phys. **B277** (1986) 456;
A. Vainshtein, V. Zakharov and M. Shifman, Sov. J. Nucl. Phys. 43 (1986) 1028;
M. Shifman, A. Vainshtein and V. Zakharov Phys. Lett. **B166** (1986) 334
- [3] J. Ellis, M. Karliner and M.A. Samuel, hep-ph/9612202
- [4] T. van Ritbergen, J.A.M. Vermaseren and S.A. Larin, hep-ph/9701390
- [5] P.M. Ferreira, I. Jack, D.R.T. Jones and C.G. North hep-ph/9705328
- [6] S.G. Gorishny, A.L. Kataev, S.A. Larin and L.R. Surguladze, Phys. Lett. **B256** (1991) 81
- [7] P.M. Ferreira, I. Jack, D.R.T. Jones, hep-ph/9702304